

Electroweak Baryogenesis driven by an **Axion-Like Particle**

Chang Sub Shin (IBS-CTPU)

*Based on work [arXiv:1806.02591, 1811.03294 [hep-ph]]
with Kwang Sik Jeong (PNU) and Tae Hyun Jung (IBS-CTPU)*


at YuCHE 2019

Feb. 27, 2019

Idea of EWBG


Electroweak baryogenesis (EWBG)

Baryon asymmetry of the Universe should be answered by physics beyond the SM. However, the Higgs still can play an important role to trigger EWBG by first order electroweak phase transition (EWPT).




Electroweak baryogenesis (EWBG)


Baryon asymmetry of the Universe should be answered by physics beyond the SM. However, the Higgs still can play an important role to trigger EWBG by first order electroweak phase transition (EWPT).



EWBG 1 (2)



EWBG 2 (2)




Extension for first order phase transition

Most of extensions beyond the SM focuses on realizing strong 1st order EWPT


(single field description)

(multi field description)


I. Thermally (BEC) Driven





IIA. Tree-Level (Ren.) Driven



IIB. Tree-Level (Non-Ren.) Driven



III. Loop Driven



[Chung, Long, Wang 12]


Extension for first order phase transition

Most of extensions beyond the SM to realize strong 1st order EWPT *needs strong couplings*

(single field description)


(multi field description)

I. Thermally (BEC) Driven




light scalars with large coupling to the Higgs without VEV

IIA. Tree-Level (Ren.) Driven




singlet scalars with cubic coupling to the Higgs, sH^+H

IIB. Tree-Level (Non-Ren.) Driven




Strong dynamics to give a low cutoff

III. Loop Driven



(charged) light fermions with large couplings to the Higgs



multi step PT with strong couplings (for Z_2 symmetric case: [Kurup, Perelstein 17])

[Chung-Leng, Wang 12]

**Landau
Pole**

Flavor

**Dark
Matter**

LHC search

**Higgs
precision**

$$\frac{n_B}{s} = 0.8 \times 10^{-10}$$

EWBG

EDM

**Higgs
mass**

**hierarchy
problems**

GW

Landau Pole

Flavor

D

LHC search

Run II @ 13 TeV

Higgs precision

$$\frac{n_B}{s} = 0.8 \times$$

EWBG

EDM

ACME II for eEDM

Higgs mass

hierarchy problems

g

1. Naturally safe from EDM and LHC constraints

Successful strong 1st order EWPT
and sizable CPVs for EWBG
with only feeble interactions?

$$\frac{n_B}{s} = 0.8 \times 10^{-10}$$


EWBG

LHC search

EDM


2. New experimental searches for the evidence of EWBG?

With Axion-Like Particles (ALPs)



2. New experimental searches for the evidence of EWBG!

ALP landscapes



Theoretical motivations of ALP for various ranges of its mass and decay constant

Axionic EWBG

[Jeong, Jung, CSS 18]

Outline

ALP intro

- compact, light, suppressed by a large axion decay constant

Strong first order phase transition

- With only feeble interactions *independently from a decay constant*

Generation of baryon asymmetry

- Non-local, local electroweak baryogenesis *depending on a decay constant*

CP violating sources

- Dynamical Top Yukawa, Electroweak theta term


Experimental constraints

- Natural suppression of EDM.
- ALP searches (LHC, meson rare decays, Supernova cooling)

ALP intro (1)

ALP, $a(x)$, is the scalar field in effective theories well below the scale f :

1) The SM singlet, and compact with a period: $2\pi f$




$$S[a] = S[a + 2\pi\mathbb{N}f]$$

ALP intro (2)

ALP, $a(x)$, is the scalar field in effective theories well below the scale f :

2) Approximate continuous shift symmetry $U(1)_{PQ}$

($a \rightarrow a + 2\pi f\beta$, where $\beta \in \mathbb{R}$)



The potentials and interactions to explicitly break shift symmetry are generated at a scale (μ) much lower than f ($\mu \ll f$)

ALP intro (3)

All interactions between ALP and matters can be given by the combination of

$$\frac{a}{f}$$

A natural way to introduce *higher dim. operators*, *weak couplings*, and *small mass of ALP*

E.g. for $\Lambda \ll f$

$$V(a) = -\Lambda^4 \cos \frac{a}{f} = \Lambda^4 + \frac{\Lambda^4}{2f^2} a^2 - \frac{\Lambda^4}{24f^4} a^4 + \frac{\Lambda^4}{720f^6} a^6 + \dots$$

the ALP mass,

$$m_a = \frac{\Lambda^2}{f} \ll f, \Lambda$$

the self coupling,

$$\lambda_{\text{quartic}} = \frac{\Lambda^4}{6f^4} \ll O(1)$$

Axion couplings to matters

$$\mathcal{L} \ni \frac{a}{16\pi^2 f} (c_G G \tilde{G} + c_W W \tilde{W} + c_B B \tilde{B}) + x_q e^{ia/f} H Q_L q_R + h.c. + \dots$$

The energy scale that we concern: $E \ll f \rightarrow \Gamma_{\text{int}} \propto (E/f)^2$

Axionic extension of the Higgs potential

A scalar potential is constructed by the Higgs and the angular field, $\theta(x) \equiv a(x)/f$

$$V(H, a) = V(H^+ H, \sin \theta, \cos \theta).$$

As an simple example with $\mu_1 \sim \mu_2 \sim \Lambda \sim O(m_W)$ (a UV model will is presented later)

$$V(H, a) = \mu_1^2 |H|^2 + \lambda |H|^4 + \mu_2^2 \cos(\theta + \alpha) |H|^2 - \Lambda^4 \cos \theta.$$

Considering an expansion in terms of a/f ,

$$V(h, a) = \frac{1}{2} \left(\mu^2 + c_1 \frac{\mu^2}{f} a + c_2 \frac{\mu^2}{f^2} a^2 + c_3 \frac{\mu^2}{f^3} a^3 + \dots \right) h^2 + \frac{\lambda}{4} h^4 \\ + \frac{\Lambda^4}{2f^2} a^2 - \frac{\Lambda^4}{24f^4} a^4 + \frac{\Lambda^4}{720f^6} a^6 + \dots.$$


The couplings between ALP and the Higgs are suppressed for $m_W \ll f$.

Tadpole, cubics and higher dimensional operators can be systematically introduced without worrying about stability of the scalar potential.

A strong 1st order EWPT can be realized!


Schematic description of the potential

The scalar potential can be written as $V(h, \theta) = \tilde{V}(\theta) + \frac{1}{2}m^2(\theta)h^2 + \frac{\lambda}{4}h^4$.



Schematic description of the potential


The scalar potential can be written as $V(h, \theta) = \tilde{V}(\theta) + \frac{1}{2}m^2(\theta)h^2 + \frac{\lambda}{4}h^4$.



The potential is bounded from below due to the periodicity of the axion dependence

Schematic description of EWPT


The scalar potential can be written as $V_T(h, \theta) = \tilde{V}(\theta) + \frac{1}{2}(m^2(\theta) + cT^2)h^2 + \frac{\lambda}{4}h^4$ for a large value of f ($T \leq m_W \ll f$), since the axion is not thermalized. at $0 < T < T_c$




Schematic description of EWPT

The scalar potential can be written as $V_T(h, \theta) = \tilde{V}(\theta) + \frac{1}{2}(m^2(\theta) + cT^2)h^2 + \frac{\lambda}{4}h^4$

at $T_c < T$



Schematic description of EWPT




Strong first order EWPT

After fixing parameters by the Higgs mass and the Higgs VEV from (with $\mu_1^2 > 0, \mu_2^2 < 0$)

$$V_{tree}(h, a) = \frac{\mu_1^2}{2} h^2 + \frac{\lambda}{4} h^4 + \frac{\mu_2^2}{2} \cos(\theta + \alpha) h^2 - \Lambda^4 \cos \theta$$

the free parameters are

$$\Lambda, \alpha, \epsilon = \sqrt{2\lambda}\Lambda^2 / (-\mu_2^2) \quad (\text{not } f)$$




Non-local generation of baryon asymmetry

Most of baryons are generated at symmetric phase after CP violating diffusion

$$\frac{dn_B}{dt} + 3Hn_B = \frac{\Gamma_{sph}(\text{sym})}{T} \left(\mu_B(\text{sym}) - c_0 \frac{n_B}{T^2} \right)$$

chemical potential
Induced by diffusion





Local generation of baryon asymmetry

Adiabatically generated inside the bubble wall by Higgs dependent chemical potential

$$\frac{dn_B}{dt} + 3Hn_B = \frac{\Gamma_{sph}(h)}{T} \left(\mu_B(h) - c_0 \frac{n_B}{T^2} \right)$$





*non-zero chemical potential
inside a bubble wall*



f determines the bubble wall width

For large $f \gg m_W$,
$$L_W \sim \frac{f}{m_W^2} \sim \frac{1}{T_n} \left(\frac{f}{m_W} \right)$$

$f \lesssim O(10 - 100) \text{ TeV} : \text{non-local gen.}$
 $f \gtrsim O(10 - 100) \text{ TeV} : \text{local gen.}$



CP violation

CP violating source (which depends on the bubble wall profile) from

$$\mathcal{L}_{CPV} \ni \frac{a}{f} O_{SM}(x)$$

During phase transition ($\Delta h \sim O(m_W)$), $\Delta\theta = \Delta a/f \sim O(1)$ even for a very large f
: Enhancing CPV effects

Examples:

1) *Dynamical top Yukawa coupling [1806.02591]*

$$Y_t(\theta) h t_L t_R + h.c., \quad \text{where } Y_t(\theta) = (y_t + x_t e^{i\theta})$$

2) *Dynamical electroweak theta-term [1811.03294] ($\mu_B \propto \dot{\theta}$)*


$$\frac{g_2^2}{16\pi^2} \Theta(\theta) \text{Tr}[W_{\mu\nu} \tilde{W}^{\mu\nu}], \quad \text{where } \Theta(\theta) = \theta$$

$$(\partial_\mu J_B^\mu) = \frac{N_f}{8\pi^2} (W_{\mu\nu} \tilde{W}^{\mu\nu} - B_{\mu\nu} \tilde{B}^{\mu\nu})$$

An UV example (1)

$$(y_t + x_t e^{i\theta}) h t_L t_R + h.c.$$

As a UV model, we can propose that the PQ symmetry is anomalously broken by hidden sector confining gauge symmetry.



$$\mathcal{L}_{eff}^{(1)} = -m_N N\bar{N} + \frac{yy'}{m_L} N\bar{N} |H|^2 + \frac{\kappa N\bar{N}}{m_S^2 m_T} H Q_L t_R + h.c.$$


$N + \bar{N}$ hidden quarks, condensate: $\langle N\bar{N} \rangle = \Lambda_h^3 e^{i\alpha(x)/f}$ from axion-hidden meson mixing

$$\mathcal{L}_{eff}^{(2)} = -m_N \Lambda_h^3 \cos \theta + \frac{yy' \Lambda_h^3}{m_L} \cos(\theta + \alpha) |H|^2 + \frac{\kappa \Lambda_h^3}{m_S^2 m_T} e^{i(\theta + \beta)} H Q_L t_R + h.c.$$

An UV example (2)

$$(y_t + x_t e^{i\theta}) h t_L t_R + h.c.$$

As a UV model, we can propose that the PQ symmetry is anomalously broken by hidden sector confining gauge symmetry.



$$\mathcal{L}_{eff}^{(1)} = -m_N N\bar{N} + \frac{yy'}{m_L} N\bar{N} |H|^2 + \frac{\kappa N\bar{N}}{m_S^2 m_T} H Q_L t_R + h.c.$$


$N + \bar{N}$ hidden quarks, condensate: $\langle N\bar{N} \rangle = \Lambda_h^3 e^{i a(x)/f}$ from axion-hidden meson mixing

$$\mathcal{L}_{eff}^{(2)} = -m_N \Lambda_h^3 \cos \theta + \frac{yy' \Lambda_h^3}{m_L} \cos(\theta + \alpha) |H|^2 + \frac{\kappa \Lambda_h^3}{m_S^2 m_T} e^{i(\theta + \beta)} H Q_L t_R + h.c.$$


Results

For $f \sim O(1 - 10)$ TeV $(y_t + x_t e^{i\theta}) h t_L t_R$

EDM and collider constraints are easily evaded for a sufficiently large value of f



EDM $\propto 1/f^2$



$$\mathcal{L}_{eff} = -\frac{\mu_0^4}{2} \frac{(\delta a)^2}{f^2} + \mu_1^3 \frac{\delta a}{f} \delta h + \mu_2^3 \frac{(\delta a)^2}{f^2} \delta h + i \frac{\delta a}{f} x_t m_t \bar{t} \gamma_5 t + h.c.$$

New! $|d_e| < 1.1 \times 10^{-29}$ cm \cdot e at 90% C.L


$m_a \sim m_w \left(\frac{m_w}{f} \right) \rightarrow m_a \sim O(0.1 - 10)$ GeV

ALP searches

After integrating out top and Higgs, ALP couplings to <gluon, photon, light quark and lepton> are generated. *Model dependent (axion decay channels) constraints are applied*


$$\mathcal{L}_{eff} = \frac{1}{16\pi^2} \frac{(\delta a)}{f} (c_g G_{\mu\nu} \tilde{G}^{\mu\nu} + c_\gamma F_{\mu\nu} \tilde{F}^{\mu\nu} + \dots) + \frac{\delta a}{f} \delta_{mix} m_q \bar{q}q + \frac{\delta a}{f} \delta_{mix} m_\ell \bar{\ell}\ell$$

Axion with mass around (5 – 10)GeV is model independently safe.



[Flacke, Frugiuele, Fuchs, Gupta, Perez 16]

[Choi, Im 16]





[Jaeckel, Spannowsky 15]

ALP searches

After integrating out top and Higgs, ALP couplings to <gluon, photon, light quark and lepton> are generated. *Model dependent (axion decay channels) constraints are applied*

$$\mathcal{L}_{eff} = \frac{1}{16\pi^2} \frac{(\delta a)}{f} \left(c_g G_{\mu\nu} \tilde{G}^{\mu\nu} + c_\gamma F_{\mu\nu} \tilde{F}^{\mu\nu} + \dots \right) + \frac{\delta a}{f} \delta_{\text{mix}} m_q \bar{q}q + \frac{\delta a}{f} \delta_{\text{mix}} m_\ell \bar{\ell}\ell$$

Axion with mass around (5 – 10) GeV is model independently safe.



[Knapen, Lin, Lou, Melia 17]



ALP searches

After integrating out top and Higgs, ALP couplings to <gluon, photon, light quark and lepton> are generated. *Model dependent (axion decay channels) constraints are applied*

$$\mathcal{L}_{eff} = \frac{1}{16\pi^2} \frac{(\delta a)}{f} \left(c_g G_{\mu\nu} \tilde{G}^{\mu\nu} + c_\gamma F_{\mu\nu} \tilde{F}^{\mu\nu} + \dots \right) + \frac{\delta a}{f} \delta_{\text{mix}} m_q \bar{q}q + \frac{\delta a}{f} \delta_{\text{mix}} m_\ell \bar{\ell}\ell$$

Axion with mass around (5 – 10)GeV is model independently safe.



Interesting hints for $m_a \sim 10 - 20$ GeV



[Mariotti, Redigolo, Sala, Tobioka 17]

For $f > O(10 - 100) \text{ TeV}$ $\frac{g_2^2 \theta}{16\pi^2} \text{Tr}[W_{\mu\nu} \tilde{W}^{\mu\nu}]$

Baryon asymmetry is nearly independent of f for $f < O(10^7 \text{ GeV})$




Totally safe from EDM and LHC constraints!

ALP searches

There is the interesting allowed window for $f \sim 10^6 - 10^7$ GeV ($m_a \sim 5 - 100$ MeV)

$\delta = \text{axion-Higgs mixing} \propto 1/f$



Conclusions

- *Axionic extension of the Higgs potential gives new parameter spaces for singlet extensions of EWBG: weakly coupled, controllable higher dimensional operators.*
- *EWPT and its cosmological evolution show different features compared to usual EWBS models: We can get stronger first order phase transition to compensate large bubble wall effects.*
- *Non-local and local baryogenesis can be realized depending on the axion decay constant.*
- *Interesting mass ranges of the ALP mass between 5-20 GeV and 5-100MeV are motivated by EWBG for the target of future ALP searches.*